SOLBIOTETM TREHALOSE SG

CHb

Bringing Sustainable Solutions to Biopharmaceuticals

Our high-purity and low-endotoxin saccharide products provide you solutions for biologics, vaccines, and cell-based therapeutics development.

Discover our philosophy: Pharmaceutical Ingredients x Sustainability

> PLATINUM Top 1% ecovadis Sustainability Rating MAR 2024



Achieving equitable well-being for all; no one shall be left behind

We are dedicated to the sustainable production of life-changing excipients, SOLBIOTE[™], to create a prosperous future for both people and the planet. We aspire to go beyond simply providing pharmaceutical solutions to deliver universal health; we ensure equitable access to quality healthcare for all.

Product inquiries to:

Y=mx+6

5

m.a

Nagase & Co., Ltd.

Life & Healthcare Products Department dnfct@ex.nagase.co.jp





SOLBIOTE[™]

Injectable Grade of Maltose

General

Maltose is a reducing disaccharide consisting of two glucose molecules linked by an α -1,4 bond. Maltose is manufactured from starch using enzyme technology.

Historically maltose has been a common disaccharide consumed by humans when assimilated into the body from consuming carbohydrate based foods. More recently, maltose has been used as an excipient for tablets and granulation, and as a carbon source in cell culture medium.


MALTOSE PH is highly purified crystalline maltose monohydrate, and has low endotoxin.

MALTOSE PH is intended for injection and intravenous fluid as energy source etc.

Chemical formula: $C_{12}H_{22}O_{11} \cdot H_2O$ Molecular weight: 360.31 CAS RN®: 6363-53-7

Properties

- High purity maltose
- Low endotoxin level guaranteed
- Good water solubility (61.9 g/100 g-H₂0 \cdot 20°C)
- Relatively compatible with oils
- The surface tension of a 1% solution is almost the same value as pure water.
- Because it is highly purified, maltose can be used for injection, as an excipient for solid, inhalation and topical formulations, and as a reagent.



SOLBIOTE[™]

Scientific reports

Maltose has been studied and reported also in various non-pharmaceutical applications.

Purpose	Title	Reference
Growth inducer of culture embryo	Breeding of new citrus cultivars by Embryo culture (No.1) Effect of saccharides on embryo growth <i>in vitro</i> .	Bulletin of the Agricultural Research Institute of Kanagawa Prefecture,133, 75-81 (1991)
Growth inducer of culture plant microspore	Formation of Calli from Isolated Microspore Cultures of Asiatic Hybrid Lily 'Connecticut King'	Journal of the Japanese Society for Horticultural Science, 69(1), 52-56 (2000)
Growth inducer of culture leaf disc	Effects of Sugar on callus and organ formation from leaf segments of Chrysanthemum	Plant Tissue Culture Letters, 3(2), 71-77 (1986)
Growth inducer of culture wheat anther	An efficient anther culture method for obtaining a higher frequency of pollen embryos in <i>Triticum aestivum</i> L.	Bullentin of the RIAR, Ishikawa Agricultural College, 3, 19-25 (1993)
Growth inducer of culture rice anther	Improvement of medium and the use in rice anther culture	Bull. Tokushima. Pref. Agri. Res. Ins., 3, 31-38 (2006)
Improver of productivity of actives from culture cells	Intermittent maltose feeding enhances pacilitaxel production in suspension culture of Taxus chinensis cells.	Biotechnol Lett., 22, 22, 1793-1796 (2000)

Packaging

25 kg (PE bag in carton box)

MANUFACTURER : Nagase Viita Co., Ltd. CONTACT : Nagase & Co., Ltd.

Life & Healthcare Products Department E-mail: dnfct@ex.nagase.co.jp

The information provided herein is intended only for reference purposes. It is the customer's responsibility to determine that the ingredients meet all legal requirements in the country where they are used, and that they do not infringe on any third party patents. Unauthorized reproduction of this brochure is prohibited.

\land Nagase Viita